

# A Unique Overlap of Scarring and Non-scarring Alopecia in Primary Cutis Verticis Gyrata

© Batuhan Mustafa Demir<sup>1</sup>, © Serkan Yazici<sup>1</sup>, © Şaduman Balaban Adım<sup>2</sup>

<sup>1</sup>Department of Dermatology and Venereology, Bursa Uludağ University Faculty of Medicine, Bursa, Türkiye

<sup>2</sup>Department of Pathology, Bursa Uludağ University Faculty of Medicine, Bursa, Türkiye

## Abstract

Cutis verticis gyrata (CVG) is a rare scalp disorder characterized by cerebriform thickening and folding of the scalp skin. It can present as a primary essential form without underlying systemic involvement. Alopecia, classified as scarring or non-scarring, may rarely accompany CVG. We report a 55-year-old female with primary essential CVG exhibiting both non-scarring alopecia areata and scarring lichen planopilaris, which were confirmed histopathologically. No neurological, ophthalmological, or systemic abnormalities were present. Magnetic resonance imaging revealed no intracranial or cranial bone pathology. To our knowledge, this is the first reported case of simultaneous scarring and non-scarring alopecia in a patient with primary essential CVG. This rare coexistence may reflect the presence of multiple autoimmune pathways targeting different segments of the hair follicle. Clinicians should consider combined pathological mechanisms in unusual alopecia presentations associated with CVG.

**Keywords:** Cutis verticis gyrata, alopecia areata, lichen planopilaris, scarring alopecia, autoimmune overlap

## INTRODUCTION

Cutis verticis gyrata (CVG) is a rare dermatological condition characterized by excessive thickening and folding of the scalp skin, resulting in a cerebriform appearance that mimics the surface of the brain. CVG is classified into three subtypes: primary essential, primary non-essential (associated with neuropsychiatric or ophthalmological disorders), and secondary (due to local or systemic causes such as tumors, inflammatory dermatoses, or trauma).<sup>1</sup> The pathogenesis of primary CVG remains poorly understood. Proposed mechanisms include hormonal imbalances, such as altered androgen metabolism, and abnormalities in connective tissue development. Histopathological findings in primary CVG are often non-specific, ranging from normal skin to hypertrophy of adnexal structures and increased dermal collagen. Alopecia is broadly divided into non-scarring and scarring types, depending on whether permanent destruction of hair follicles occurs. Alopecia areata (AA) is a common autoimmune form

of non-scarring alopecia, whereas lichen planopilaris (LPP) represents a lymphocytic scarring alopecia.<sup>2</sup> To date, only a few cases have been reported in the literature that associate alopecia with CVG, and none have described the coexistence of both scarring and non-scarring types in the same patient with primary essential CVG. This report presents a unique case illustrating this rare overlap.

## CASE REPORT

A 55-year-old woman presented with progressive scalp deformity and hair loss. Ten years earlier, she had been diagnosed with seborrheic dermatitis and CVG based on scalp biopsy, but no follow-up was performed. On examination, alopecic patches and deep cerebriform folds were observed over the vertex and occipital scalp. Histopathology from these areas showed features of both scarring and non-scarring

Submission: 04-May-2025

Epub: 14-Aug-2025

Acceptance: 14-Jul-2025

### Access this article online

Quick Response Code:



Website:

www.turkjdermatol.com

DOI:

10.4274/tjd.galenos.2025.54376

**Address for correspondence:** Batuhan Mustafa Demir, MD,

Department of Dermatology and Venereology, Bursa Uludağ University

Faculty of Medicine, Bursa, Türkiye


Email: batuhanmdemir@uludag.edu.tr

ORCID ID: 0009-0005-3276-2813



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given.

**How to cite this article:** Demir BM, Yazici S, Adım ŞB. A unique overlap of scarring and non-scarring alopecia in primary cutis verticis gyrata. Turk J Dermatol. 2025;19(3):118-121.



**Figure 1.** Histopathologic sections from scalp biopsies: (A) Follicular hypertrophy in dermis (H&E x 40). (B) Follicular scar tissue (H&E x 100). (C) Perifollicular lichenoid infiltrate at infundibular level, suggestive of LPP (H&E x 200). (D) Mixed pattern: LPP below, AA above in same field (H&E x 100). (E) Lymphocytic infiltration around anagen hair bulb (H&E x 200).  
H&E: Hematoxylin and eosin, LPP: Lichen planopilaris, AA: Alopecia areata



**Figure 2.** The clinical image of the patient's scalp shows the CVG pattern in the occipital region  
CVG: *Cutis verticis gyrata*

alopecia: perifollicular lichenoid inflammation and fibrosis consistent with LPP, along with peribulbar lymphocytic infiltrates characteristic of alopecia areata (Figure 1). Systemic evaluation, including cranial magnetic resonance imaging, revealed no neurologic or ocular involvement or intracranial pathology (Figure 2). She was started on topical corticosteroids and scheduled for regular dermatologic follow-up. Written informed consent was obtained from the patient for publication.

## Discussion

CVG is an uncommon condition involving hypertrophy of the scalp skin, forming undulating folds that resemble cerebral gyri. It may present as a primary essential form, with no systemic associations, or secondary to underlying local or systemic conditions.<sup>1,3</sup> Histopathology in primary CVG is often non-specific, though thickening of dermal collagen or adnexal hypertrophy may be observed. In our case, histologic features of CVG coexisted with findings diagnostic for both LPP (scarring alopecia) and AA (non-scarring alopecia). Although

**Table 1. Cases of CVG with alopecia**

Literature	Age	Gender	CVG type	Alopecia type
Buontempo et al. <sup>7</sup>	39	F	Primary essential	Primary scarring alopecia
Anansiripun and Suchonwanit <sup>8</sup>	24	M	Secondary	Scarring alopecia
Alonso Pereira et al. <sup>9</sup>	29	M	Secondary	Folliculitis decalvans, folliculitis keloidalis nuchae
Bonalumi Filho et al. <sup>10</sup>	43	F	Secondary	N/A
Alcántara González et al. <sup>11</sup>	48	M	Secondary	N/A
Yoo et al. <sup>6</sup>	28	M	Primary essential	Alopecia areata
Mishra et al. <sup>12</sup>	18	F	Primer essential	N/A
Saoji et al. <sup>13</sup>	48	M	Secondary	N/A
Pai and Rao <sup>14</sup>	25	M	Secondary	Scarring alopecia
Fox et al. <sup>15</sup>	Newborn	F	Congenital	N/A
van Geest et al. <sup>16</sup>	46	M	Secondary	N/A
Jeanfils et al. <sup>17</sup>	43	M	Secondary	N/A
Hamm and Argent <sup>18</sup>	41	F	Secondary	N/A
Yazici et al. <sup>19</sup>	7	F	Secondary	N/A
Kanwar et al. <sup>20</sup>	25	F	N/A	Tractional

N/A: No answer, F: Female, M: Male, CVG: Cutis verticis gyrata

both LPP and AA are considered autoimmune disorders, they typically exhibit distinct immunopathological profiles. The co-occurrence of scarring and non-scarring alopecia in the same patient raises the possibility of a shared underlying autoimmune predisposition, potentially targeting different components or layers of the hair follicle unit. This aligns with theories proposing that variable follicular immune privilege collapse may contribute to divergent clinical outcomes.<sup>4,5</sup> A literature review identified 15 prior cases of CVG with alopecia. Of these, one involved AA<sup>6</sup> and one primary CVG with scarring alopecia (Table 1).<sup>7</sup> However, to our knowledge, this is the first reported case of both forms of alopecia occurring simultaneously in a patient with primary essential CVG. Further studies are needed to clarify whether this dual pattern reflects coincidental findings or a broader, multi-level immune dysregulation affecting follicular structures in CVG patients.

## Ethics

**Informed Consent:** Written informed consent was obtained from the patient for publication.

## Footnotes

## Authorship Contributions

Design: S.Y., Analysis or Interpretation: Ş.B., Literature Search: B.M.D., Writing: B.M.D.

**Conflict of Interest:** The authors declared that they have no conflict of interest.

**Financial Disclosure:** The authors declared that this study received no financial support.

## REFERENCES

- Chamli A, Jones M, Bacha T, Litaïem N, Zeglaoui F. Cutis verticis gyrata: three cases illustrating three different etiologies. *Clin Case Rep*. 2022;10(5):e05889.
- Al Aboud AM, Syed HA, Zito PM. Alopecia. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2024.
- Palazzo R, Schepis C, Ruggeri M, Baldini L, Pizzimenti A, Arcoraci V, Spina E. An endocrinological study of patients with primary cutis verticis gyrata. *Acta Derm Venereol*. 1993;73(5):348-349.
- Mofarrah R, Mofarrah R, Jahani Amiri K, Ghasemi M. Co-localization of alopecia areata and lichen planopilaris in a patient receiving immunosuppressants: a rare case. *J Cosmet Dermatol*. 2021;20(3):976-979.
- Mohaghegh F, Bahrami B, Saber M. The simultaneous occurrence of lichen planopilaris and alopecia areata: a report of two cases and literature review. *Clin Case Rep*. 2020;8(9):1622-1627.
- Yoo KH, Lee JW, Jang WS, Li K, Seo SJ, Hong CK. Cutis verticis gyrata and alopecia areata: a synchronous coincidence? *Yonsei Med J*. 2010;51(4):612-614.
- Buontempo MG, Alhanshali L, Shapiro J, Klein EJ, Oh CS, Kim RH, Rodriguez EA, Lo Sicco K. A case of cutis verticis gyrata developing in a patient with primary scarring alopecia: a unique presentation of a rare disorder. *JAAD Case Rep*. 2023;38:44-47.
- Anansiripun P, Suchonwanit P. A case of progressive thickening and furrowing of facial skin and scalp with scarring alopecia. *Skin Appendage Disord*. 2021;7(3):237-242. doi: 10.1159/000513792.
- Alonso Pereira L, Teixeira M, Andrade J, Rochael M, Gavazzoni-Dias MFR. An overlap of secondary cutis verticis gyrata, folliculitis decalvans, folliculitis keloidalis nuchae and the use of dreadlocks: the role of inflammation due to traction. *Skin Appendage Disord*. 2017;2(3-4):130-134.
- Bonalumi Filho A, Darrigo LG Jr, Avelaira JC, Kac BK, Azulay DR. Nevo intradérmico cerebriforme como causa de cutis verticis gyrata [Intradermal cerebriform nevus as cause of cutis verticis gyrata]. *Rev Assoc Med Bras* (1992). 2010;56(6):639-641.

11. Alcántara González J, Truchuelo Díez MT, Carrillo Gijón R, Martín Díaz RM, Jaén Olasolo P. Letter: Cerebriform intradermal nevus presenting as secondary cutis verticis gyrata. *Dermatol Online J*. 2010;16(12):14.
12. Mishra A, Tehrani H, Hancock K, Duncan C. Management of primary cutis verticis gyrata with tissue expansion and hairline lowering foreheadplasty. *J Plast Reconstr Aesthet Surg*. 2010;63(6):1060-1061.
13. Saoji V, Chaudhari S, Gohokar D. Primary systemic amyloidosis: three different presentations. *Indian J Dermatol Venereol Leprol*. 2009;75(4):394-397.
14. Pai VG, Rao GS. Congenital cerebriform melanocytic naevus with cutis verticis gyrata. *Indian J Dermatol Venereol Leprol*. 2002;68(6):367-368.
15. Fox LP, Geyer AS, Anyane-Yeboah K, Garzon MC. Cutis verticis gyrata in a patient with Noonan syndrome. *Pediatr Dermatol*. 2005;22(2):142-146.
16. van Geest AJ, Berretty PJ, Klinkhamer PJ, Neumann HA. Cerebriform intradermal naevus (a rare form of secondary cutis verticis gyrata). *J Eur Acad Dermatol Venereol*. 2002;16(5):529-531.
17. Jeanfils S, Tennstedt D, Lachapelle JM. Cerebriform intradermal nevus. a clinical pattern resembling cutis verticis gyrata. *Dermatology*. 1993;186(4):294-297.
18. Hamm JC, Argenta LC. Giant cerebriform intradermal nevus. *Ann Plast Surg*. 1987;19(1):84-88.
19. Yazici AC, Ikizoglu G, Baz K, Polat A, Ustunsoy D. Cerebriform intradermal nevus. *Pediatr Dermatol*. 2007;24(2):141-143.
20. Kanwar AJ, Ghosh S, Thami GP, Kaur S. Alopecia and cutis verticis gyrata due to traction presenting as headache. *Int J Dermatol*. 1992;31(9):671-672.